

BIOME GUIDE PROTOCOL

Allergies & Hay Fever Microbiome Protocol

Evidence-based prebiotic, probiotic, and dietary strategies
to rebalance immune tolerance through the gut-lung axis

Covers: Seasonal Allergic Rhinitis | Perennial Allergic Rhinitis | Hay Fever | Atopic Allergy

Based on 15 systematic reviews, meta-analyses, and RCTs from PubMed | March 2026

biomeguide.com

What This Protocol Addresses

Allergic rhinitis (AR) affects roughly 10-30% of the global population and is the most common allergic condition worldwide. Whether you suffer from seasonal hay fever triggered by pollen or year-round symptoms from dust mites and pet dander, the underlying immunological pattern is the same: your immune system has lost tolerance to harmless environmental substances and mounts an inappropriate Th2-dominant inflammatory response, producing excess IgE antibodies that trigger mast cell degranulation, histamine release, and the sneezing, itching, congestion, and watery eyes you know too well.

Groundbreaking research has revealed that the gut microbiome is a central regulator of immune tolerance. Through the gut-lung axis, microbial metabolites and immune cells traffic between your gut and respiratory tissues, directly influencing whether your immune system reacts to pollen as a threat or tolerates it as harmless. This protocol targets the microbial imbalance that underlies allergic disease: restoring depleted SCFA-producing bacteria, rebalancing the Th1/Th2 immune axis, and providing natural mast cell-stabilizing support to reduce your allergic burden from the inside out.

Timeline expectation: For seasonal allergies, begin this protocol 4 weeks before your allergy season for best results. Probiotic effects on immune rebalancing take 4-8 weeks to reach full benefit based on RCT timelines. Quercetin and butterbur provide faster symptomatic relief (within days to weeks). Full microbial remodeling takes 3-6 months of consistent adherence. For perennial AR, follow the protocol year-round with seasonal adjustments.

The Gut-Allergy Connection

Your gut houses approximately 70% of your immune system. The relationship between gut microbiota and allergic disease operates through the gut-lung axis, a bidirectional communication network linking intestinal immune regulation to respiratory health.

- 1. Th1/Th2 Imbalance:** Healthy gut microbiota promotes balanced immune responses. In allergic individuals, microbial dysbiosis allows Th2 dominance, driving IgE production, mast cell activation, and eosinophilic inflammation. Reduced SCFA-producing bacteria impair regulatory T cell (Treg) differentiation, removing the brake on allergic Th2 responses.
- 2. SCFA Depletion:** AR patients consistently show reduced fecal butyrate levels (approximately 30-40% lower than controls). Butyrate has a negative correlation with serum IgE levels and regulates nasal mucosal barriers. Propionate and butyrate inhibit histone deacetylase activity, promoting Treg differentiation and suppressing the Th2 responses that drive allergic symptoms (Yang et al., 2025; Ke et al., 2025).
- 3. Early-Life Origins:** Delayed gut microbiota maturation in the first year of life is a hallmark of pediatric allergic disease. Children who develop allergies harbor less mature microbiomes with lower

Bifidobacterium and *Clostridia* abundance during infancy (Hoskinson et al., 2023).

What the meta-analyses show: Luo et al. (2022) analyzed 28 RCTs and found probiotics significantly improved quality of life in AR (RQLQ: SMD -0.64, $p < 0.001$) and reduced symptoms (SMD -0.29, $p < 0.001$). Xi et al. (2025) found a 25% reduced risk of allergic disease with probiotics (RR 0.75, $p = 0.009$). Lu et al. (2025) ranked probiotic regimens across 31 RCTs (2,544 patients), finding mixed formulations most effective overall.

Sources: Yang et al. (2025) Front Microbiol. DOI: 10.3389/fmicb.2025.1654997; Ke et al. (2025) Curr Res Microb Sci. DOI: 10.1016/j.crmicr.2025.100362; Luo et al. (2022) Front Immunol. DOI: 10.3389/fimmu.2022.848279; Xi et al. (2025) Front Nutr. DOI: 10.3389/fnut.2025.1502390

The Allergy Microbial Fingerprint

Research across multiple meta-analyses and systematic reviews reveals a consistent pattern of microbial dysbiosis in allergic individuals: depletion of protective, SCFA-producing bacteria and enrichment of pro-inflammatory species. Understanding this fingerprint is the foundation for targeted restoration.

Depleted taxa (targets for restoration):

Bacterium	Role	Evidence
<i>Bifidobacterium</i> spp. (esp. <i>B. longum</i>)	Immune regulation, Treg promotion, SCFA production, Th1/Th2 balancing	Consistently depleted in AR, food allergy, and atopic children
<i>Faecalibacterium prausnitzii</i>	Major butyrate producer, anti-inflammatory, inversely correlated with fecal IgE	Reduced in AR patients; negatively correlated with IgE
<i>Lactobacillus</i> spp.	Lactic acid production, immune modulation, barrier support	Depleted in allergic disease broadly
<i>Roseburia</i> spp.	Butyrate producer, gut barrier maintenance	Depleted; reduced SCFA output in AR
<i>Clostridium butyricum</i>	Butyrate producer; reduces serum IgE by ~22% in studies	Depleted in AR (Yang et al., 2025)
<i>Dorea</i> spp.	Negatively correlated with fecal IgE in dust mite AR	Reduced in AR (Ke et al., 2025)

Enriched taxa (pro-inflammatory, elevated in allergy):

Bacterium	Concern
<i>Ruminococcus gnavus</i>	Mucin degrader; consistently elevated in allergic children
<i>Escherichia coli</i>	LPS producer; 4.6-fold elevation in wheezing/atopic children
Enterobacteriaceae	LPS-producing family; enriched in allergic conditions
<i>Collinsella</i> spp.	Increases intestinal permeability; enriched in atopic disease
Proteobacteria	LPS-producing phylum; increased in AR patients

Sources: Li et al. (2025) PeerJ. DOI: 10.7717/peerj.19441; Ke et al. (2025) Curr Res Microb Sci. DOI: 10.1016/j.crmicr.2025.100362; Hoskinson et al. (2023) Nat Commun. DOI: 10.1038/s41467-023-40336-4

What to AVOID

Histamine-rich foods (during peak symptoms): Aged cheeses, wine (especially red), fermented meats, and sauerkraut contain high histamine levels that compound your body's allergic histamine response. Paradoxically, some fermented foods that benefit gut health can worsen symptoms during peak allergy season.

Cross-reactive foods (Oral Allergy Syndrome): If you have birch pollen allergy, raw apples, cherries, and peaches may trigger oral itching. Ragweed: melons, bananas. Mugwort: celery, carrots. Cooking denatures the cross-reactive proteins and usually eliminates the reaction.

Processed foods and emulsifiers: Carboxymethylcellulose (CMC) and polysorbate-80 damage your intestinal mucus layer and promote the dysbiosis that underlies allergic disease.

Refined sugar: Promotes inflammatory Th2 responses and feeds pro-inflammatory Proteobacteria. High sugar intake is associated with increased allergic sensitization.

Excessive alcohol: Histamine content (especially in wine and beer) and vasodilating effects worsen nasal congestion. Alcohol also increases intestinal permeability.

Dairy (individual assessment): May thicken mucus and worsen congestion in some individuals during acute allergic episodes. Response varies; try a 2-week elimination during peak season to assess personal sensitivity.

Dietary Recommendations

Anti-Allergic Mediterranean Framework

The Mediterranean diet pattern naturally provides the three pillars of dietary allergy management: polyphenols (natural mast cell stabilizers), prebiotic fibers (to restore depleted gut flora), and omega-3 fatty acids (anti-inflammatory).

Quercetin-rich foods (natural antihistamine): Onions (especially red onions), apples, berries, capers, green tea, and citrus fruits. Quercetin inhibits mast cell degranulation and histamine release. **Prebiotic-rich foods:** Garlic, leeks, asparagus (inulin/FOS sources), oats, green bananas, cooled rice and potatoes (resistant starch). **Omega-3 foods:** Fatty fish (salmon, sardines, mackerel), walnuts, flaxseed.

Seasonal Strategy

Pre-season (4 weeks before): Maximize prebiotic intake and begin probiotic supplementation to prime your immune system before pollen exposure peaks. This timing aligns with the 4-week window needed for probiotics to influence immune parameters.

During season: Limit histamine-rich foods; avoid cross-reactive raw fruits/vegetables (cook them instead); increase quercetin-rich foods; maintain consistent prebiotic and probiotic intake.

Off-season: Focus on long-term microbiome restoration; introduce fermented foods (kefir, yogurt, kimchi) when histamine burden is lower; continue prebiotic diversity.

Supplement Protocol

Prebiotic Supplements (Foundation)

These prebiotic fibers feed your depleted beneficial bacteria. The prebiotic approach restores your own endogenous SCFA-producing flora, which in turn promotes Treg cells and suppresses the Th2 responses that drive your allergic symptoms. Start with half doses for the first week.

Prebiotic	Feeds	Allergy-Specific Notes	Dose
Inulin / FOS	<i>Bifidobacterium</i> (strong), <i>Faecalibacterium</i>	Restores the most consistently depleted genus in allergy	Start 1/2 tsp, titrate to 1 tsp
GOS	<i>Bifidobacterium</i> (strongest), <i>Lactobacillus</i>	Highly selective bifidogenic; IgE-lowering potential	3.5g/day
Resistant Starch	<i>Roseburia</i> , <i>Faecalibacterium</i> , <i>R. bromii</i>	Butyrate production for Treg support and IgE reduction	1 heaping Tbsp/day
Apple Pectin	<i>Faecalibacterium</i> , <i>Eubacterium</i>	Gentle fermentation; good for sensitive individuals	1 heaping Tbsp/day
Arabinoxylans	<i>Bifidobacterium</i> , <i>Roseburia</i>	Bifidogenic; supports SCFA production	1 heaping Tbsp/day
Polyphenols	<i>Akkermansia muciniphila</i>	Dual benefit: Akkermansia support + direct mast cell stabilization	1/2 tsp berry powder

Probiotic Recommendations

A network meta-analysis of 31 RCTs (2,544 AR patients) found that mixed probiotic formulations demonstrated the best combined efficacy for reducing nasal symptoms, improving quality of life, and lowering IgE levels (Lu et al., 2025). Several specific strains also have individual RCT support.

Strain	Evidence	Dose	Duration
<i>B. longum</i> BB536 + <i>B. infantis</i> M-63 + <i>B. breve</i> M-16V	RCT (n=40 children, 8 wks): Symptom scores dropped from 9.3 to 3.5 (p<0.005) while placebo worsened. (Del Giudice et al., 2017)	BB536: 3×10^9 + M-63: 1×10^9 + M-16V: 1×10^9 CFU/day	8+ weeks
<i>L. paracasei</i> LP-33	GA2LEN RCT (n=425, 5 wks): Significant RQLQ improvement (p=0.0255); significant ocular symptom relief (p=0.0029). (Costa et al., 2014)	Per product label	5+ weeks; through allergy season

Strain	Evidence	Dose	Duration
<i>B. longum</i> BB536 (single)	RCT (n=40, 14 wks): Significantly alleviated eye symptoms (OR 0.31, p=0.044); tended to relieve nasal symptoms. (Xiao et al., 2006)	5x10 ¹⁰ CFU/day	4-14 weeks
Multi-strain mix (highest ranked)	Network meta-analysis of 31 RCTs: mixed formulations ranked best for TNSS, RQLQ, and IgE reduction. (Lu et al., 2025)	Per product label	8+ weeks

Choosing your strains: The Bifidobacterium triple combination (BB536 + M-63 + M-16V) has the strongest individual RCT evidence for allergic rhinitis. If you are already taking antihistamines, LP-33 showed benefit as an add-on therapy. For maximum coverage, choose a multi-strain product containing both *Bifidobacterium* and *Lactobacillus* species. **Start 4 weeks before allergy season** for best preventive effect.

Supporting Supplements

Supplement	Why	Dose	Duration
Quercetin	Natural mast cell stabilizer. Meta-analysis of 13 RCTs (823 pts): significantly reduced TNSS, sneezing, and nasal itching in seasonal AR. Inhibits histamine release without drowsiness.	500-1,000 mg/day (phytosomal form for absorption)	During allergy season; or year-round for perennial AR
Vitamin D3	Meta-analysis of 5 RCTs: significant benefit when combined with non-corticosteroid treatment (SMD -0.56, $p < 0.001$). Promotes Tregs and Th1/Th2 balance. (Kawada et al., 2025)	2,000-5,000 IU/day (test levels; target 40-60 ng/mL)	Ongoing; retest at 3 months
Butterbur (Ze339)	RCT (n=125): As effective as cetirizine for seasonal AR, with fewer sedative side effects. Six RCTs support efficacy. MUST use PA-free extract only. (Schapowal et al., 2002)	8mg petasines, 4x/day	During acute seasonal symptoms only
Sodium Butyrate	Directly addresses core SCFA depletion; butyrate negatively correlated with IgE; promotes Treg cells; regulates nasal mucosal barriers.	300-600 mg/day microencapsulated	First 60 days
Vitamin C	Natural antihistamine properties; supports immune function; may reduce severity of allergic symptoms.	1,000-2,000 mg/day	During allergy season

Safety notes: Butterbur must be a PA-free (pyrrolizidine alkaloid-free) extract; raw butterbur is hepatotoxic. Do not use butterbur if you are allergic to ragweed, daisies, or marigolds (same plant family). Test vitamin D levels before high-dose supplementation. Quercetin may interact with certain antibiotics and blood thinners; consult your provider. Always discuss new supplements with your healthcare provider, especially if you take prescription allergy medications or immunotherapy.

The Daily Protocol

The Daily Prebiotic Shake

The Daily Prebiotic Shake

1. Add at least 12 oz of water to a shaker bottle or blender
2. Add 1 heaping tablespoon of stabilized rice bran (arabinoxylans)
3. Add 1 heaping tablespoon of apple pectin powder
4. Add 1 heaping tablespoon of resistant starch (potato starch or Hi-Maize 260)
5. Add 1 heaping teaspoon of inulin powder (start with 1/2 tsp week 1)
6. Add 1/2 teaspoon berry polyphenol powder (mast cell support + *Akkermansia*)
7. Shake well and drink!

Timing: Once daily off-season; twice daily during allergy season. **Week 1:** Use HALF of all prebiotic doses. Increase to full dose by week 2-3 as tolerated.

The Two-Phase Protocol

Phase 1: Active Immune Rebalancing (4 weeks pre-season through peak season, or Weeks 1-8 for perennial AR)

During Phase 1, you are actively rebuilding protective gut flora, rebalancing the Th1/Th2 axis, and providing direct anti-allergic support. Start this phase 4 weeks before your allergy season begins.

- Daily prebiotic shake (once daily; increase to twice daily during peak symptoms)
- Probiotic: Bifidobacterium triple mix or multi-strain, daily
- Quercetin: 500-1,000 mg/day (phytosomal form, with meals)
- Vitamin D3: 5,000 IU/day with a fat-containing meal
- Sodium butyrate: 300-600 mg/day with meals
- Vitamin C: 1,000-2,000 mg/day
- Butterbur (Ze339): during acute symptoms only (PA-free extract)
- Follow anti-allergic Mediterranean diet; limit histamine-rich foods during peak season
- Avoid cross-reactive raw fruits/vegetables (cook them instead)
- Avoid processed foods, refined sugar, emulsifiers, and excessive alcohol

Phase 2: Maintenance (Off-season / Year-round for perennial AR)

In Phase 2, you maintain your microbial gains and prepare your immune system for the next season. For perennial AR, Phase 2 is your ongoing baseline with Phase 1 intensity during flare-ups.

- Reduce prebiotic shake to once daily or every other day
 - Continue probiotics (can rotate strains every 2-3 months)
 - Reduce quercetin to 250-500 mg/day maintenance dose
 - Continue vitamin D3 (retest at 3 months; adjust to maintenance 2,000 IU/day)
 - Discontinue sodium butyrate (your own bacteria should now produce more)
 - Discontinue butterbur (seasonal use only)
 - Introduce fermented foods (kefir, yogurt, kimchi) when histamine burden is lower
 - Begin next season's probiotic protocol 4 weeks before pollen season starts
 - Focus on quercetin-rich whole foods: red onions, apples, berries, green tea
-

What to Expect

Timeframe	What You May Experience
Week 1	Adjustment period. Mild increase in gas or bloating as your microbiome shifts. Quercetin and vitamin C may provide early symptomatic relief. Butterbur (if used) provides relief within days, comparable to cetirizine.
Weeks 2-4	Bloating resolves. Probiotics begin influencing immune parameters. If started pre-season, your immune system is being primed for better tolerance when pollen arrives. Nasal congestion and sneezing may begin to improve.
Weeks 4-8	Full probiotic benefit window. Based on RCT timelines, this is when quality of life scores (RQLQ) and nasal symptom scores (TNSS) show meaningful improvement. Reduced reliance on rescue antihistamines.
Months 2-3	Sustained immune rebalancing. Th1/Th2 ratio normalization. Reduced frequency and severity of allergic episodes. Less reactivity to previously problematic triggers. Transition to maintenance phase.
Months 3-6	Full microbial remodeling. Measurable increases in <i>Bifidobacterium</i> , <i>Faecalibacterium</i> , and overall diversity on microbiome testing. Many patients report their "best allergy season ever" after consistent protocol adherence.

Tracking Your Progress

Keep a simple daily symptom journal during allergy season: rate sneezing, congestion, itching, and eye symptoms each on a 0-3 scale (Total Nasal Symptom Score). Track rescue antihistamine use. Compare season-over-season to measure improvement. The RQLQ (Rhinoconjunctivitis Quality of Life Questionnaire) is a validated tool your allergist can administer for objective tracking.

Consider a microbiome test (such as Biomesight, Ombre, or Thryve) at baseline and after 3-6 months to objectively measure changes in your *Bifidobacterium*, *Faecalibacterium*, and overall diversity. Tracking IgE levels with your allergist provides another objective marker of improvement.

Learn more at BiomeGuide.com

Every probiotic strain and prebiotic mentioned in this protocol has a detailed profile page on our site with links to the original research. Visit biomeguide.com to explore our probiotic strain database, condition-specific guides, and microbiome testing resources.

Sources

Every recommendation in this protocol is based on published, peer-reviewed clinical research. Below is the complete citation list with DOIs for verification.

1. Li M, et al. (2025) "Association between gut microbiota and allergic rhinitis: a systematic review and meta-analysis." *PeerJ*, 13:e19441. DOI: 10.7717/peerj.19441
2. Yang W, et al. (2025) "Gut-lung axis in allergic rhinitis: microbial dysbiosis and therapeutic strategies." *Front Microbiol*, 16:1654997. DOI: 10.3389/fmicb.2025.1654997
3. Xi Z, et al. (2025) "Efficacy of probiotics in the treatment of allergic diseases: a meta-analysis." *Front Nutr*, 12:1502390. DOI: 10.3389/fnut.2025.1502390
4. Luo C, et al. (2022) "The efficacy and safety of probiotics for allergic rhinitis: a systematic review and meta-analysis." *Front Immunol*, 13:848279. DOI: 10.3389/fimmu.2022.848279
5. Ke H, et al. (2025) "Advances in research on gut microbiota and allergic diseases in children." *Curr Res Microb Sci*, 8:100362. DOI: 10.1016/j.crmicr.2025.100362
6. Lu C, et al. (2025) "Efficacy of different probiotic regimens for allergic rhinitis: A network meta-analysis." *Complement Ther Clin Pract*, 59:101954. DOI: 10.1016/j.ctcp.2025.101954
7. Costa DJ, et al. (2014) "Efficacy and safety of the probiotic *Lactobacillus paracasei* LP-33 in allergic rhinitis: GA2LEN study." *Eur J Clin Nutr*, 68(5):602-7. DOI: 10.1038/ejcn.2014.13
8. Xiao JZ, et al. (2006) "Effect of probiotic *Bifidobacterium longum* BB536 in relieving symptoms of Japanese cedar pollinosis." *J Investig Allergol Clin Immunol*, 16(2):86-93. PMID: 16689181
9. Xiao JZ, et al. (2007) "Clinical efficacy of probiotic *Bifidobacterium longum* BB536 for cedar pollen allergy." *Allergol Int*, 56(1):67-75. DOI: 10.2332/allergolint.O-06-455
10. Del Giudice MM, et al. (2017) "Bifidobacterium mixture treatment in children with seasonal AR and intermittent asthma." *Ital J Pediatr*, 43:25. DOI: 10.1186/s13052-017-0340-5
11. Kawada K, et al. (2025) "Vitamin D Supplementation and Allergic Rhinitis: A Systematic Review and Meta-Analysis." *Medicina*, 61(2):355. DOI: 10.3390/medicina61020355
12. Schapowal A, et al. (2002) "Randomised controlled trial of butterbur and cetirizine for treating seasonal allergic rhinitis." *BMJ*, 324:144. DOI: 10.1136/bmj.324.7330.144
13. Hoskinson C, et al. (2023) "Delayed gut microbiota maturation in the first year of life is a hallmark of pediatric allergic disease." *Nat Commun*, 14:4785. DOI: 10.1038/s41467-023-40336-4
14. Anandan C, et al. (2009) "Omega 3 and 6 oils for primary prevention of allergic disease: systematic review and meta-analysis." *Allergy*, 64(6):840-8. DOI: 10.1111/j.1398-9995.2009.02042.x
15. Clinical Effects of Polyphenolic Compounds on Allergic Rhinitis: A Systematic Review and Meta-Analysis. *JACI: In Practice* (2025). DOI: 10.1016/j.jaip.2025.06.013

This information is for educational purposes only and is not medical advice. Consult your healthcare provider before starting any new supplement regimen, especially if you take prescription allergy medications, are undergoing allergen immunotherapy, or have asthma. These protocols are designed to complement, not replace, your medical treatment. Individual results may vary based on allergy severity, specific allergens, and duration of symptoms. If you experience severe allergic reactions, difficulty breathing, or anaphylaxis, seek emergency medical attention immediately. BiomeGuide.com is not affiliated with any specific supplement manufacturer.

Ready to learn more?

Visit **biomeguide.com** to explore our probiotic strain database, condition guides, and microbiome testing resources. Every strain mentioned in this protocol has a detailed profile page with links to the original research.

biomeguide.com