

BIOME GUIDE PROTOCOL

Candida & Fungal Overgrowth Protocol

Evidence-based prebiotic, probiotic, and dietary
strategies to restore microbial balance and
suppress Candida overgrowth

Covers: GI Candida Overgrowth | SIFO | Recurrent Vulvovaginal Candidiasis | Fungal Dysbiosis

Based on 12 systematic reviews and RCTs from PubMed | April 2026

biomeguide.com



What This Protocol Addresses

Candida albicans is the most common fungal inhabitant of the human gastrointestinal tract, colonizing an estimated 40-80% of healthy adults. Under normal conditions, your resident bacterial microbiota keeps *Candida* in check through a set of defense mechanisms collectively called **colonization resistance**. When this balance is disrupted -- by antibiotics, immunosuppressive drugs, proton pump inhibitors (PPIs), high-sugar diets, or chronic stress -- *Candida* can shift from its harmless yeast form into an invasive hyphal (filamentous) form, leading to overgrowth and a range of symptoms.

This protocol targets the root cause of *Candida* overgrowth: **a depleted bacterial microbiota that has lost its ability to suppress fungal expansion**. Rather than relying solely on antifungal drugs (which don't prevent recurrence), this approach rebuilds the protective bacterial community that keeps *Candida* in check naturally, while simultaneously using evidence-based dietary and supplement strategies to directly inhibit *Candida* virulence factors.

The protocol covers gastrointestinal *Candida* overgrowth, Small Intestinal Fungal Overgrowth (SIFO), and crossover strategies relevant to recurrent vulvovaginal candidiasis (VVC). All recommendations are grounded in peer-reviewed clinical research.

Timeline expectation: Dietary and supplement changes can produce noticeable improvements in bloating, gas, and digestive discomfort within 2-4 weeks. Meaningful reduction in *Candida* colonization typically requires 4-8 weeks of consistent intervention. For recurrent VVC, maintenance protocols of 3-6 months are supported by RCT evidence. Full microbial rebalancing -- where your bacterial community consistently suppresses *Candida* without ongoing supplementation -- may take 3-6 months.

The Gut-Candida Connection

Your gut bacteria suppress *Candida* through multiple mechanisms. **Short-chain fatty acids (SCFAs)** -- particularly butyrate, produced by beneficial bacteria fermenting dietary fiber -- directly impair *Candida* growth, block the yeast-to-hyphae transition (the key virulence switch), and maintain the oxygen-depleted colonic environment that favors bacteria over fungi. A landmark review by Bays and Savage (2025) demonstrated that butyrate promotes oxygen consumption by colonocytes, maintaining colonic hypoxia that suppresses *Candida* proliferation. Additionally, *Bacteroides* species increase HIF-1-alpha, promoting production of the antimicrobial peptide LL-37 (cathelicidin), which has direct anti-*Candida* activity.

Source: Bays & Savage (2025) *Infection and Immunity*. DOI: 10.1128/iai.00610-24

When antibiotics, PPIs, or other disruptors reduce these protective bacteria, *Candida* seizes the opportunity. It transitions to its hyphal form, which penetrates the gut lining, forms biofilms resistant to immune clearance, and triggers inflammatory cascades. This creates a self-reinforcing cycle: *Candida* overgrowth further disrupts the bacterial community, which further reduces colonization resistance, which allows more *Candida* expansion (Musumeci et al., 2022; Li et al., 2022).

SIFO (Small Intestinal Fungal Overgrowth): Two clinical studies found SIFO in 25-26% of patients with unexplained GI symptoms (bloating, belching, nausea, diarrhea). SIFO frequently co-occurs with SIBO, and risk factors include PPI use and small intestinal dysmotility. A 2-3 week course of antifungal therapy may improve symptoms, but without addressing the underlying microbial imbalance, recurrence is common (Erdogan & Rao, 2015; Soliman et al., 2025).

The Candida Overgrowth Microbial Fingerprint

Depleted taxa (targets for restoration):

Bacterium	Role in Candida Resistance	Evidence
<i>Faecalibacterium prausnitzii</i>	Major butyrate producer; butyrate directly inhibits <i>C. albicans</i> growth and filamentation; maintains colonic hypoxia	Bays & Savage (2025); Nguyen et al. (2011)
<i>Bifidobacterium</i> spp.	Produce acetate/lactate that cross-feed butyrate producers; competitive exclusion; immune modulation	Li et al. (2022); Kumamoto et al. (2020)
<i>Lactobacillus</i> spp.	Produce lactic acid, H ₂ O ₂ , bacteriocins; directly inhibit <i>Candida</i> growth, adhesion, and biofilm	Matsubara et al. (2016); Kumar et al. (2013)
<i>Roseburia</i> spp.	Butyrate producer; contributes to colonization resistance via SCFA production	Bays & Savage (2025)
<i>Bacteroides</i> spp.	Increase HIF-1-alpha, promoting LL-37 antimicrobial peptide; direct anti- <i>Candida</i> immune activation	Bays & Savage (2025); Li et al. (2022)

Enriched / Problematic organisms:

Organism	Concern
<i>Candida albicans</i> (hyphae form)	Primary pathogenic form; expresses adhesins (HWP1), invasins; forms biofilms; penetrates epithelial barriers
Non-albicans <i>Candida</i> spp.	<i>C. glabrata</i> , <i>C. tropicalis</i> , <i>C. parapsilosis</i> increasingly common, especially with fluconazole exposure
Proteobacteria	Often co-elevated with <i>Candida</i> overgrowth, indicating broader dysbiosis and inflammation

What to AVOID

Refined sugars and high-fructose corn syrup: Glucose and fructose are Candida's primary fuel sources. *C. albicans* metabolizes simple sugars far more efficiently than complex carbohydrates. A pilot study of 120 patients found that dietary sugar restriction combined with antifungal treatment achieved 85% cure vs 42.5% for antifungal alone (Otasevic et al., 2018).

Refined wheat flour products: A clinical study found that increased consumption of purified wheat flour was significantly associated with higher GI Candida colonization (Jeziorek et al., 2019). Replace with whole grains, almond flour, or coconut flour.

Alcohol: Disrupts gut barrier integrity, feeds yeast fermentation, and impairs mucosal immune function. Eliminate completely during Phase 1.

Excess fruit juice: Concentrated fructose without fiber. Whole fruits in moderation are acceptable (the fiber slows sugar absorption and feeds beneficial bacteria).

Medications to discuss with your provider:

- **Proton pump inhibitors (PPIs):** Associated with increased SIFO risk by reducing stomach acid that normally suppresses fungal colonization (Erdogan & Rao, 2015)
- **Broad-spectrum antibiotics:** Deplete protective bacteria, creating the dysbiosis that allows Candida expansion. Discuss alternatives when possible
- **Corticosteroids:** Suppress mucosal immune defenses against Candida. Discuss dosing with your prescriber

Dietary Recommendations

Anti-Candida Foods (eat daily)

Coconut oil (1-2 tablespoons daily): Contains capric acid, caprylic acid, and lauric acid -- all with direct anti-Candida activity. In a murine model, a coconut oil-rich diet significantly reduced GI Candida colonization compared to beef tallow or soybean oil. Switching from a high-saturated-fat diet to coconut oil reduced pre-existing colonization (Gunsalus et al., 2016). Use coconut oil for cooking, add to smoothies, or take directly.

Garlic, onions, leeks, asparagus: Natural sources of inulin and FOS that selectively feed *Bifidobacterium* and *Faecalibacterium*. Garlic also contains allicin, which has direct antifungal properties. Raw garlic is most potent.

Non-starchy vegetables: Provide diverse fibers that feed the full spectrum of SCFA-producing bacteria. Broccoli, leafy greens, cauliflower, zucchini, celery, bell peppers. These should form the

foundation of your diet during Phase 1.

Polyphenol-Rich Foods (feed *Akkermansia muciniphila*)

Berries (blueberries, raspberries, cranberries), pomegranate, green and black tea, grape seeds: Polyphenols (ellagitannins, proanthocyanidins) selectively increase *Akkermansia muciniphila*, which strengthens the mucin barrier and may limit Candida translocation from the gut lumen to deeper tissues.

Saturated and Medium-Chain Fatty Acid Sources

Cheese and dairy fats: A clinical study found that higher cheese consumption was associated with negative Candida cultures, suggesting a protective effect of saturated fatty acids against fungal colonization (Jeziorek et al., 2019). Medium-chain fatty acids (MCFAs) disrupt Candida cell membranes and inhibit biofilm formation.

Supplement Protocol

Prebiotic Supplements (rebuild colonization resistance)

Prebiotic	Feeds	Product	Dose
Arabinoxylans (rice bran)	<i>Bifidobacterium</i> , <i>Roseburia</i>	Stabilized rice bran (NOW Foods)	1 heaping Tbsp
Inulin	<i>Bifidobacterium</i> , <i>Faecalibacterium</i>	Pure inulin powder	1 heaping tsp (titrate up)
Apple Pectin	<i>Faecalibacterium</i> , <i>Eubacterium</i>	Apple pectin powder	1 heaping Tbsp
Resistant Starch	<i>Roseburia</i> , <i>Eubacterium rectale</i> , <i>Faecalibacterium</i>	Hi-Maize 260 or Bob's Red Mill Potato Starch	1 heaping Tbsp

Important: Prebiotics do NOT feed Candida. *C. albicans* primarily metabolizes simple sugars (glucose, fructose) and cannot efficiently ferment complex polysaccharides like inulin or resistant starch. The prebiotic strategy selectively advantages bacteria over fungi, boosting SCFA production that directly suppresses Candida.

Probiotic Recommendations

Unlike blanket "don't take probiotics" advice, we recommend specific strains where clinical trial evidence supports their anti-Candida activity. The prebiotic protocol is the foundation; these probiotics provide additional targeted support.

Strain(s)	Evidence	Dose	Duration
<i>Saccharomyces boulardii</i> CNCM I-745	Secretes capric acid that inhibits <i>C. albicans</i> filamentation, adhesion, and biofilm. Downregulates virulence genes HWP1, INO1, CSH1 (Murzyn et al., 2010)	250-500 mg (5-10 billion CFU) 2x daily	4-8 weeks (Phase 1), then maintenance
<i>L. rhamnosus</i> GR-1 + <i>L. reuteri</i> RC-14	RCT: With fluconazole, reduced yeast detection from 38.5% to 10.3% at 4 weeks ($p=0.014$); reduced symptomatic discharge from 34.6% to 10.3% ($p=0.03$) (Martinez et al., 2009)	1-2 billion CFU each, oral daily	4+ weeks

Strain(s)	Evidence	Dose	Duration
<i>L. acidophilus</i> GLA-14 + <i>L. rhamnosus</i> HN001 + Lactoferrin	RCT: Reduced RVVC recurrence from 100% (placebo) to 29.2% at 6 months. Significant improvement in itching and discharge (Russo et al., 2019)	Combination capsule with 100 mg lactoferrin	5-day induction, then 10 days/month for 6 months
Multi-strain: <i>L. acidophilus</i> , <i>L. rhamnosus</i> , <i>B. longum</i> , <i>B. bifidum</i> , <i>S. boulardii</i>	RCT: Reduced Candida colonization by 37.2% at day 14 (31.3% vs 50%, RR 0.63, p=0.02); candiduria reduced from 37.3% to 17.3% (p=0.006) (Kumar et al., 2013)	10+ billion CFU combined	14+ days

Supporting Supplements

Supplement	Why	Dose	Duration
Sodium Butyrate	Directly inhibits <i>C. albicans</i> growth, blocks yeast-to-hyphae transition, disrupts biofilms, enhances macrophage killing via ROS/NO (Nguyen et al., 2011)	600 mg - 2 g/day (non-delayed release)	First 60 days
Caprylic Acid	Medium-chain fatty acid; disrupts Candida cell membrane; synergistic with butyrate. Found naturally in coconut oil (Murzyn et al., 2010; Gunsalus et al., 2016)	500-1000 mg with meals, 2-3x daily	4-8 weeks (Phase 1)
Bovine Lactoferrin	Iron-binding glycoprotein; sequesters iron from Candida (essential for fungal growth); synergistic with probiotics (Russo et al., 2019)	100-200 mg/day	3-6 months
Vitamin D3	Promotes LL-37 antimicrobial peptide production; modulates mucosal immunity; deficiency associated with increased Candida susceptibility	5,000 IU/day with meal	Ongoing (test levels)
Vitamin A (Retinol)	Supports mucosal immune integrity, IgA production, gut barrier function at Candida colonization sites	10,000-25,000 IU/day	30-60 days only

Safety notes: Vitamin A at doses above 10,000 IU is **contraindicated in pregnancy** (teratogenic risk). Do not exceed 25,000 IU/day and limit to 30-60 days. Test liver function if using >10,000 IU. Vitamin D: test serum levels at 3-6 months, target 50-70 ng/mL. If you are taking immunosuppressive medications or antifungal drugs, consult your healthcare provider before starting this supplement protocol.

The Daily Protocol

The Daily Prebiotic Shake

The Daily Prebiotic Shake

1. Add at least 12 oz of water to a shaker bottle or blender
2. Add 1 heaping tablespoon of stabilized rice bran (arabinoxylans)
3. Add 1 heaping tablespoon of apple pectin powder
4. Add 1 heaping tablespoon of resistant starch (Hi-Maize 260 or potato starch)
5. Add 1 heaping teaspoon of inulin powder (titrate up over 2 weeks)
6. Add 1 teaspoon of polyphenol powder (pomegranate or cranberry) -- optional
7. Add 1 tablespoon of coconut oil (melt first if solid)
8. Shake well and drink!

Timing: Once daily, morning or midday. Take *S. boulardii* with the shake; take caprylic acid with a separate meal.

Phase 1: Active Anti-Candida Intervention (Weeks 1-6)

- Prebiotic shake once daily
- *Saccharomyces boulardii* 250-500 mg twice daily
- Caprylic acid 500-1000 mg with meals, 2-3 times daily
- Sodium butyrate 600 mg - 2 g/day
- Bovine lactoferrin 100-200 mg/day
- Vitamin D3 5,000 IU/day with fat-containing meal
- Vitamin A 10,000 IU/day (skip if pregnant or planning pregnancy)
- 1-2 Tbsp coconut oil daily (cooking or smoothie)
- Strict sugar and refined flour elimination
- No alcohol
- Avoid fermented foods temporarily (may aggravate symptoms initially in sensitive individuals)
- Prebiotic-rich whole foods daily: garlic, onion, leeks, asparagus, non-starchy vegetables

Phase 2: Maintenance & Rebalancing (Week 7 onward)

- Continue prebiotic shake (can reduce to 5x/week)
 - Reduce *S. boulardii* to 250 mg once daily (maintenance dose)
-

- Discontinue caprylic acid (or reduce to 1x daily if symptoms persist)
 - Continue sodium butyrate through day 60, then discontinue
 - Discontinue vitamin A
 - Continue vitamin D3 indefinitely (test levels at 3-6 months)
 - Continue lactoferrin for 3-6 months total
 - Cautiously reintroduce fermented foods: start with small amounts of sauerkraut, kefir
 - Gradually reintroduce moderate whole grains and natural sugars (observe for symptom return)
 - Continue cooking with coconut oil
 - Consider adding *L. rhamnosus* GR-1 / *L. reuteri* RC-14 for ongoing protection
-

What to Expect

Timeframe	What You May Experience
Weeks 1-2	Adjustment period. Mild bloating or gas may increase temporarily as prebiotic fibers shift microbial populations. Sugar cravings may intensify before diminishing. This is normal and typically resolves within 7-10 days.
Weeks 2-4	Bloating and gas improve. Digestive discomfort begins to decrease. Energy levels may improve as Candida-driven inflammation reduces. Sugar cravings diminish noticeably.
Weeks 4-8	Meaningful symptom improvement. Reduced bloating, more regular digestion, clearer skin (if Candida was contributing to skin symptoms). VVC recurrence rate drops significantly with probiotic maintenance.
Months 2-4	Microbial rebalancing becomes more established. SCFA production increases as beneficial bacteria repopulate. Candida is suppressed by restored colonization resistance rather than active intervention.
Months 4-6+	Maintenance phase. If the prebiotic/dietary approach is sustained, the bacterial community should maintain Candida suppression independently. Can gradually reduce supplement dosing while monitoring for symptom return.

Tracking Your Progress

- **Symptom journal:** Track bloating, gas, brain fog, sugar cravings, skin changes, and energy levels weekly. Look for trends over 4-8 week blocks, not day-to-day fluctuations.
- **Stool consistency:** Use the Bristol Stool Chart. Aim for Type 3-4 (well-formed, easy to pass) as the bacterial community normalizes.
- **For VVC:** Track recurrence frequency. The Russo et al. (2019) RCT showed recurrence dropped from 100% to 29.2% at 6 months with probiotic + lactoferrin maintenance.
- **Microbiome testing (optional):** A stool mycobiome + bacteriome test at baseline and at 3-6 months can show shifts in Candida levels and bacterial diversity. Look for increased *Faecalibacterium*, *Bifidobacterium*, and *Roseburia*, and decreased Candida relative abundance.
- **SIFO evaluation:** If SIFO was diagnosed by your provider, a follow-up small bowel aspirate at 8-12 weeks can confirm reduction in fungal load.

Learn more at BiomeGuide.com

Visit our probiotic strain database to explore detailed profiles for *S. boulardii*, *L. rhamnosus* GR-1, *L. reuteri* RC-14, and every other strain mentioned in this protocol. Each profile links directly to the original PubMed-indexed research.

Sources

Every recommendation in this protocol is based on published, peer-reviewed clinical research.

1. Bays HM, Savage DC. (2025) "Candida albicans gastrointestinal colonization resistance: a host-microbiome balancing act." *Infection and Immunity*. DOI: 10.1128/iai.00610-24. PMID: 40788152
2. Jawhara S. (2022) "How Gut Bacterial Dysbiosis Can Promote Candida albicans Overgrowth during Colonic Inflammation." *Microorganisms*, 10(5):1014. DOI: 10.3390/microorganisms10051014. PMID: 35630457
3. Kumamoto CA, Gresnigt MS, Hube B. (2020) "The gut, the bad and the harmless: Candida albicans as a commensal and opportunistic pathogen in the intestine." *Current Opinion in Microbiology*, 56:7-15. DOI: 10.1016/j.mib.2020.05.006. PMID: 32604030
4. Otasevic S, et al. (2018) "The dietary modification and treatment of intestinal Candida overgrowth -- a pilot study." *Journal de Mycologie Medicale*, 28(4):623-627. DOI: 10.1016/j.mycmed.2018.08.002. PMID: 30166063
5. Nguyen LN, et al. (2011) "Sodium butyrate inhibits pathogenic yeast growth and enhances the functions of macrophages." *Journal of Antimicrobial Chemotherapy*, 66(11):2573-2580. DOI: 10.1093/jac/dkr358. PMID: 21911344
6. Matsubara VH, et al. (2016) "Probiotics as Antifungals in Mucosal Candidiasis." *Clinical Infectious Diseases*, 62(9):1143-1153. DOI: 10.1093/cid/ciw038. PMID: 26826375
7. Kumar S, et al. (2013) "Evaluation of efficacy of probiotics in prevention of Candida colonization in a PICU -- a randomized controlled trial." *Critical Care Medicine*, 41(2):565-572. DOI: 10.1097/CCM.0b013e31826a409c. PMID: 23361033
8. Li XV, et al. (2022) "Interactions between Candida albicans and the resident microbiota." *Frontiers in Microbiology*, 13:930495. DOI: 10.3389/fmicb.2022.930495. PMID: 36204612
9. Murzyn A, et al. (2010) "Capric acid secreted by *S. boulardii* inhibits *C. albicans* filamentous growth, adhesion and biofilm formation." *PLoS ONE*, 5(8):e12050. DOI: 10.1371/journal.pone.0012050. PMID: 20706577
10. Martinez RC, et al. (2009) "Improved treatment of vulvovaginal candidiasis with fluconazole plus probiotic *Lactobacillus rhamnosus* GR-1 and *Lactobacillus reuteri* RC-14." *Letters in Applied Microbiology*, 48(3):269-274. DOI: 10.1111/j.1472-765X.2008.02477.x. PMID: 19187507
11. Russo R, et al. (2019) "Randomised clinical trial in women with Recurrent Vulvovaginal Candidiasis: Efficacy of probiotics and lactoferrin as maintenance treatment." *Mycoses*, 62(4):328-335. DOI: 10.1111/myc.12883. PMID: 30565745
12. Gunsalus KTW, et al. (2016) "Manipulation of Host Diet To Reduce Gastrointestinal Colonization by the Opportunistic Pathogen *Candida albicans*." *mSphere*, 1(1):e00020-15. DOI: 10.1128/mSphere.00020-15. PMID: 27303684
13. Erdogan A, Rao SSC. (2015) "Small intestinal fungal overgrowth." *Current Gastroenterology Reports*, 17(4):16. DOI: 10.1007/s11894-015-0436-2. PMID: 25786900
14. Soliman N, et al. (2025) "Small Intestinal Bacterial and Fungal Overgrowth: Health Implications and Management Perspectives." *Nutrients*, 17(8):1365. DOI: 10.3390/nu17081365. PMID: 40284229
15. Jeziorek M, et al. (2019) "The influence of diet on gastrointestinal *Candida* spp. colonization and the susceptibility of *Candida* spp. to antifungal drugs." *Roczniki Panstwowego Zakladu Higieny*, 70(2):195-200. DOI: 10.32394/rpzh.2019.0070. PMID: 31215785
16. Musumeci S, et al. (2022) "The human gut mycobiome and the specific role of *Candida albicans*: where do we stand, as clinicians?" *Clinical Microbiology and Infection*, 28(1):58-63. DOI: 10.1016/j.cmi.2021.07.034. PMID: 34363944

This information is for educational purposes only and is not medical advice. Consult your healthcare provider before starting any new supplement regimen, especially if you are taking antifungal medications, immunosuppressive drugs, or are pregnant or planning pregnancy. Individual results may vary. BiomeGuide.com is not affiliated with any specific supplement manufacturer.

Ready to learn more?

Visit biomeguide.com to explore our probiotic strain database, condition guides, and microbiome testing resources. Every strain mentioned in this protocol has a detailed profile page with links to the original PubMed-indexed research.

biomeguide.com
