

BIOME GUIDE PROTOCOL

Common Cold & URTI Microbiome Protocol

Evidence-based prebiotic, probiotic, and dietary strategies to strengthen immune resilience through the gut-lung axis

Covers: Common Cold | Viral URTI | Seasonal Respiratory Illness | Immune Resilience

Based on 16 clinical studies and systematic reviews from PubMed | April 2026

biomeguide.com

What This Protocol Addresses

The common cold is the most frequent infectious disease in humans. Adults average 2-4 colds per year, while children can catch 6-10. Most are caused by rhinoviruses, though coronaviruses, adenoviruses, and influenza viruses also contribute to upper respiratory tract infections (URTIs). Each episode typically lasts 7-10 days, with symptoms peaking around days 2-3.

Conventional advice focuses on hand washing, rest, and symptom management after you are already sick. This protocol takes a fundamentally different approach: it targets the gut-lung axis to strengthen your immune system's ability to prevent and shorten respiratory infections before they take hold. Your gut microbiome is not just about digestion -- it is the command center for roughly 70% of your immune system.

A Cochrane review of 23 randomized controlled trials (6,950 participants) found that probiotics reduced the number of people diagnosed with URTIs by 24% (RR 0.76), reduced URTI incidence rates by 18% (rate ratio 0.82), and cut antibiotic prescriptions by 42% (RR 0.58). Prebiotic supplementation showed similar benefits, reducing subjects with at least one RTI by 27% (OR 0.73). These are meaningful, clinically significant reductions backed by thousands of participants (Zhao et al., 2022; Williams et al., 2022).

Timeline expectation: Start this protocol 4-6 weeks before cold and flu season for best results. Immune priming through the gut-lung axis takes time. You may notice subtle improvements in how you feel within 2-3 weeks (fewer minor throat irritations, faster recovery from minor bugs). By 6-8 weeks, your gut microbiome should be measurably shifted toward higher SCFA production and enhanced NK cell activity. The real payoff is over the full cold season: fewer colds, shorter duration, and less severe symptoms.

The Gut-Lung Axis

The gut-lung axis is the bidirectional communication pathway between your gastrointestinal and respiratory systems. When your gut bacteria ferment dietary fiber, they produce short-chain fatty acids (SCFAs) -- primarily butyrate, propionate, and acetate. These SCFAs travel through your bloodstream to the lungs, where they directly enhance the function of alveolar macrophages (your lungs' first-line defense), dendritic cells (which coordinate the immune response), and natural killer (NK) cells (which destroy virus-infected cells) (Chunxi et al., 2020).

This is not theoretical. A systematic review found that during respiratory tract infections, gut microbiome diversity (Shannon index) decreased by 1.45 units compared to healthy controls ($p < 0.0001$). Key SCFA-producing bacteria -- Lachnospiraceae, Ruminococcaceae, and Ruminococcus -- were consistently depleted, while the opportunistic pathogen Enterococcus was enriched (Woodall et al., 2022). The gut-lung connection is real, measurable, and modifiable.

Sources: Chunxi et al. (2020) J Immunol Res; Woodall et al. (2022) PLoS One; Zhao et al. (2022) Cochrane Database Syst Rev

The URTI Microbial Fingerprint

Research has identified consistent gut microbiome differences between people who are susceptible to frequent respiratory infections and those who resist them. The pattern centers on bacteria that produce immune-modulating short-chain fatty acids.

Depleted Taxa (targets for restoration):

Bacterium	Role	Evidence
Lachnospiraceae (family)	Major SCFA producers; butyrate and propionate production; immune modulation via gut-lung axis	Woodall et al. (2022): depleted in RTI patients vs healthy controls
Ruminococcaceae (family)	Butyrate producers; fiber fermentation; immune homeostasis maintenance	Woodall et al. (2022): depleted in RTI patients
Ruminococcus spp.	Primary fiber degraders; keystone genus for resistant starch fermentation	Woodall et al. (2022): depleted in RTI patients
Bifidobacterium spp.	Immune modulation; IgA stimulation; acetate/lactate cross-feeding to butyrate producers	Williams et al. (2022): prebiotic targeting of Bifidobacterium reduced RTI incidence
Faecalibacterium prausnitzii	Major butyrate producer; anti-inflammatory; gut barrier and gut-lung axis support	Depleted across inflammatory conditions; supports SCFA supply to lungs

Enriched / Potentially Problematic Taxa:

Bacterium	Concern
Enterococcus	Opportunistic; pro-inflammatory; associated with gut dysbiosis during RTIs (Woodall et al., 2022)

Sources: Woodall et al. (2022) PLoS One; Williams et al. (2022) Adv Nutr

What to Avoid

Immune resilience depends on maintaining a healthy gut microbiome. Certain dietary and lifestyle factors actively undermine the gut-lung axis and increase your susceptibility to respiratory infections. Managing these is as important as adding beneficial supplements.

Dietary factors that weaken immune resilience:

Excess refined sugar: High sugar intake suppresses neutrophil and macrophage function for hours after consumption. It also feeds opportunistic bacteria (*Enterococcus*, *Proteobacteria*) at the expense of SCFA-producing *Lachnospiraceae* and *Bifidobacterium*.

Ultra-processed foods: Emulsifiers, artificial sweeteners, and food additives disrupt the mucus layer and reduce microbiome diversity. Lower diversity correlates directly with increased RTI susceptibility (Woodall et al., 2022).

Excessive alcohol: Disrupts gut barrier integrity ("leaky gut"), reduces secretory IgA in the airways, and depletes SCFA-producing bacteria. Even moderate drinking during cold season increases infection risk.

Low-fiber diets: Without fermentable fiber, your gut bacteria cannot produce the SCFAs that fuel the gut-lung axis. The Western diet (high fat, low fiber) is associated with reduced *Lachnospiraceae* and *Ruminococcaceae* -- exactly the taxa depleted during RTIs.

Discuss with your doctor: Proton pump inhibitors (PPIs) alter gut microbiome composition and have been associated with increased pneumonia risk. Frequent or unnecessary antibiotic use devastates SCFA-producing bacteria. If you need antibiotics, restart the prebiotic protocol immediately afterward to rebuild your gut flora.

Lifestyle factors: Chronic sleep deprivation (<6 hours/night) reduces NK cell activity by up to 70%. Chronic psychological stress elevates cortisol, which suppresses mucosal immunity and shifts the gut microbiome toward dysbiosis. Overtraining without adequate recovery increases URTI risk in athletes -- the "open window" hypothesis (Lagowska & Bajerska, 2021). These factors compound, so addressing them alongside the supplement protocol amplifies your results.

Sources: Woodall et al. (2022) PLoS One; Lagowska & Bajerska (2021) J Athl Train

Dietary Recommendations

Prebiotic-Rich Foods (Eat Daily)

These foods provide the fermentable fibers that feed SCFA-producing gut bacteria, fueling the gut-lung immune axis. Consistency matters more than quantity -- eat these every day, not just occasionally.

- **Onions, garlic, leeks, asparagus:** Natural sources of inulin and FOS that selectively feed Bifidobacterium and support cross-feeding to butyrate producers. Garlic also has direct antimicrobial and immunomodulatory properties.
- **Oats and barley:** Beta-glucans feed Bifidobacterium and have direct immunomodulatory effects, activating macrophages and NK cells independent of the microbiome pathway.
- **Bananas (slightly green), Jerusalem artichokes, chicory root:** Rich sources of inulin and resistant starch that feed Ruminococcus, Roseburia, and Faecalibacterium.
- **Cold potatoes and cooked-then-cooled rice:** Retrograded resistant starch that specifically feeds the Ruminococcaceae and Lachnospiraceae families depleted during RTIs.

Polyphenol-Rich Foods (Antiviral + Microbiome Support)

Polyphenols have both direct antiviral properties and support Akkermansia muciniphila, which strengthens the gut mucus barrier. Include these daily:

- **Berries (blueberries, elderberries, blackberries):** Elderberry has specific evidence for reducing cold duration and severity. All berries provide anthocyanins that support Akkermansia.
- **Green and black tea:** Catechins (EGCG) have direct antiviral activity against rhinoviruses and support microbiome diversity. 2-3 cups daily.
- **Pomegranate, dark chocolate (70%+ cacao), red/purple grapes:** Rich in ellagitannins and flavonoids that are metabolized by gut bacteria into bioactive compounds.

Fermented Foods (Live Cultures)

Several of the strongest probiotic RCTs for URTI prevention used fermented dairy as the delivery vehicle. Include daily fermented foods for complementary live cultures and postbiotics:

- **Yogurt and kefir:** Provide Lactobacillus and Bifidobacterium strains plus postbiotic metabolites. The Shida et al. (2017) trial used fermented milk (Yakult-type) with L. casei Shirota.
- **Sauerkraut, kimchi, miso:** Diverse Lactobacillus species and anti-inflammatory metabolites. Kimchi also contains garlic and ginger with direct antimicrobial properties.

Immune-Supporting Nutrients from Food

- **Vitamin C-rich foods:** Bell peppers, citrus, kiwi, broccoli, and strawberries. Regular intake reduces cold duration by 8% in adults and 14% in children (Hemila & Chalker, 2013).
- **Zinc-rich foods:** Oysters, red meat, pumpkin seeds, chickpeas, and cashews support innate and adaptive immunity.
- **Vitamin D from food:** Fatty fish (salmon, mackerel), egg yolks, and fortified foods. Most people still need supplementation, especially at higher latitudes.

Sources: Hemila & Chalker (2013) *Cochrane Database Syst Rev*; Shida et al. (2017) *Eur J Nutr*

Supplement Protocol

Prebiotic Supplements

The prebiotic protocol is the foundation of this approach. By feeding the SCFA-producing bacteria that fuel the gut-lung axis, you create sustained immune support rather than temporary stimulation. A meta-analysis of 17 studies found that prebiotic supplementation reduced the proportion of subjects experiencing at least one RTI by 27% (OR 0.73, $p=0.0002$) (Williams et al., 2022).

Prebiotic	Feeds	Product	Dose
Arabinoxylans (rice bran)	Bifidobacterium, Roseburia	Stabilized rice bran (NOW Foods)	1 heaping Tbsp
Inulin / FOS	Bifidobacterium, Faecalibacterium, Lactobacillus	Pure inulin powder	1 heaping tsp (titrate to 10g)
Apple Pectin	Faecalibacterium, Eubacterium	Apple pectin powder	1 heaping Tbsp
Resistant Starch	Roseburia, Ruminococcus, Faecalibacterium	Hi-Maize 260 or Bob's Red Mill Potato Starch	1 heaping Tbsp
GOS	Bifidobacterium (strongest bifidogenic), Lactobacillus	Bimuno GOS powder	2.5-5g/day

Probiotic Recommendations

Unlike some protocols that blanket-advise against probiotics, we recommend specific strains where randomized controlled trial evidence directly supports their use in preventing or shortening respiratory infections. The prebiotic protocol above is the foundation; probiotics provide additional, strain-specific immune support.

Strain(s)	Evidence	Dose	Duration
<i>L. paracasei</i> strain Shirota (LcS)	RCT (n=96): URTI incidence 22.4% vs 53.2% placebo ($p=0.002$). URTI-free rate significantly higher ($p=0.0008$). Maintained NK cell activity. (Shida et al., 2017)	1×10^{11} CFU/day	12+ weeks through cold season
<i>L. rhamnosus</i> GG (LGG)	Meta-analysis (4 RCTs, n=1,805 children): reduced URTIs (RR 0.62, NNT 4), otitis media (RR 0.76), antibiotic use (RR 0.80). (Liu et al., 2013)	10^9 - 10^{10} CFU/day	12+ weeks (especially for children)

Strain(s)	Evidence	Dose	Duration
B. animalis lactis BI-04	RCT (n=465): URTI risk reduced (HR 0.73, p=0.022). Median time to illness delayed 0.7 months. (West et al., 2014)	2x10 ⁹ CFU/day	5 months (full cold season)
Multi-strain: L. gasseri PA 16/8 + B. longum SP 07/3 + B. bifidum MF 20/5	RCT (n=479): shortened cold duration by ~2 days (p=0.045). Reduced severity. Increased cytotoxic T cells. (de Vrese et al., 2006)	As per product formulation	3-5 months (cold season)

Choosing your probiotic: L. paracasei Shirota (LcS) has the strongest single-strain evidence, with URTI incidence dropping from 53% to 22% in one RCT. It is commercially available as Yakult. For children, LGG has the best evidence (NNT of 4 for URTI prevention). B. lactis BI-04 is a good choice for physically active adults based on the West et al. trial. You do not need all four -- pick one based on your situation and availability.

Supporting Supplements

Supplement	Why	Dose	Duration
Vitamin D3	IPD meta-analysis (25 RCTs, 11,321 participants): reduced acute RTI risk (OR 0.88). Daily dosing: OR 0.81. In deficient subjects: 70% reduction (OR 0.30). (Martineau et al., 2017)	2,000-4,000 IU/day (daily, NOT bolus)	Year-round (test levels)
Vitamin C	Cochrane review (29 RCTs, 11,306 participants): reduced cold duration 8% in adults, 14% in children. Under physical stress: 52% incidence reduction. (Hemila & Chalker, 2013)	500-1,000mg/day	Year-round
Zinc	Cochrane review: may reduce cold duration by ~2.4 days. More useful as acute treatment (lozenges within 24h of symptom onset) than prevention. (Nault et al., 2024)	15-30mg/day preventive; 75mg/day lozenges acute	Cold season; acute: 5-7 days
Sodium Butyrate	Directly fuels the gut-lung axis by providing the key SCFA that modulates alveolar macrophage and dendritic cell function. Supports gut barrier.	300-600mg/day	Phase 1 (first 8 weeks)

Safety notes: Vitamin D at 2,000-4,000 IU/day is safe for most adults, but test your 25(OH)D levels and aim for 40-60 ng/mL. Do NOT use bolus dosing (monthly mega-doses) -- the Martineau meta-analysis showed daily dosing was effective while bolus was not. Zinc lozenges can cause nausea and taste disturbance; do not exceed 75mg/day of elemental zinc or use for more than 7 days acutely. Vitamin C doses above 2g/day may cause GI upset. If you are pregnant, breastfeeding, or on immunosuppressive medications, consult your healthcare provider before starting any supplement regimen.

The Daily Protocol

The Daily Prebiotic Shake

Your Daily Immune-Priming Shake

1. Add at least 12 oz of water to a shaker bottle or blender
2. Add 1 heaping tablespoon of stabilized rice bran (arabinoxylans)
3. Add 1 heaping tablespoon of apple pectin powder
4. Add 1 heaping tablespoon of resistant starch (Hi-Maize 260 or potato starch)
5. Add 1 heaping teaspoon of inulin powder
6. Add a scoop of elderberry or mixed berry powder (polyphenols + antiviral support)
7. Shake well and drink!

Timing: Take once daily with breakfast or lunch. During Phase 1 (immune priming), consider splitting into two half-servings if you experience GI adjustment symptoms.

The Two-Phase Protocol

Phase 1: Immune Priming (Weeks 1-8)

Ideally, start this phase 4-6 weeks before peak cold season (September-October in the Northern Hemisphere). The goal is to rebuild your SCFA-producing gut bacteria and prime the gut-lung immune axis before viral exposure peaks.

- **Weeks 1-2:** Begin prebiotic shake at lower doses (half servings). Start probiotic of choice (LcS, LGG, or BI-04). Begin vitamin D3 + vitamin C daily. Start zinc if cold season is underway. Continue eating prebiotic-rich and polyphenol-rich foods daily.
- **Weeks 3-4:** Titrate prebiotics up to full dose. Full dietary protocol. Probiotic continues. Add sodium butyrate (300-600mg/day) for direct gut-lung axis support. You may notice fewer minor throat irritations.
- **Weeks 5-8:** Full protocol. SCFA production should be increasing. NK cell activity enhanced. Gut microbiome diversity shifting. If you catch a cold during this phase, it should be shorter and milder than usual. Continue all supplements.

Phase 2: Maintenance (Through Cold Season and Beyond)

Once your gut-lung axis is primed, maintain the gains through the entire cold and flu season (typically October through April).

- **Continue prebiotic shake** once daily. You can simplify the recipe if desired, but maintain at least resistant starch + inulin + rice bran.

- **Continue probiotic** through the full cold season. After cold season ends (spring), you can cycle 1 month on / 1 month off or transition to fermented foods.
 - **Vitamin D3 + vitamin C:** Continue year-round. Test vitamin D levels annually and adjust dose to maintain 40-60 ng/mL.
 - **Zinc:** Keep zinc lozenges on hand for acute use at the first sign of a cold (within 24 hours of symptom onset). Daily preventive zinc (15mg) is optional.
 - **Discontinue sodium butyrate** after 8 weeks. Your prebiotic intake should now be generating sufficient endogenous butyrate.
 - **Continue prebiotic-rich diet** and fermented foods daily. This is a lifestyle, not just a supplement protocol.
-

Acute Cold Protocol

Even with optimal gut-lung axis support, you may still catch the occasional cold. When you feel one coming on, take immediate action within the first 24 hours for the best results.

At the First Sign of a Cold (within 24 hours of symptom onset):

1. **Zinc lozenges:** Start immediately. Use zinc acetate or zinc gluconate lozenges, totaling 75mg/day of elemental zinc. Dissolve slowly in your mouth (do not chew or swallow whole). The Cochrane review found this may reduce cold duration by ~2.4 days (Nault et al., 2024).
2. **Increase vitamin C** to 1-2g/day (split into 2-3 doses).
3. **Continue your probiotic** -- do not stop. Probiotics reduce the duration and severity of colds that do occur (de Vrese et al., 2006).
4. **Continue the prebiotic shake** -- your gut bacteria need fuel to maintain immune support.
5. **Rest and hydrate.** Reduce exercise intensity. Sleep is when your immune system does its best work.
6. **Elderberry extract** (optional): 15mL syrup or 175mg capsule, 2-4x daily for 5 days. Some evidence for reducing cold duration, though not from the gut microbiome pathway.

What to expect with the acute protocol: If you have been following the maintenance protocol and your gut-lung axis is primed, you should notice that colds are shorter (by 1-2 days) and milder (reduced congestion, sore throat, and fatigue). The combination of zinc lozenges (acute antiviral) plus an already-primed immune system (from the prebiotic/probiotic protocol) provides both immediate and sustained defense.

When to see a doctor: If symptoms worsen after day 3-4 instead of improving, if you develop a fever above 103F (39.4C), if you experience difficulty breathing, chest pain, or persistent earache, see your healthcare provider to rule out secondary bacterial infection.

For athletes and highly active individuals: A meta-analysis of 14 RCTs found that probiotic supplementation in athletes significantly reduced total symptom severity score ($p=0.02$) and lowered IL-6 ($p=0.002$) and TNF-alpha ($p=0.01$) (Lagowska & Bajerska, 2021). If you train intensely, the prebiotic + probiotic protocol is especially important during heavy training blocks and competition seasons, when the "open window" of immunosuppression is widest.

What to Expect

Timeframe	What You May Experience
Days 1-5	Possible increase in gas/bloating as prebiotics begin shifting your gut microbiome. This is normal and temporary -- it means bacteria are responding to the new fuel sources.
Weeks 1-3	Initial GI adjustment subsides. You may notice subtle improvements: fewer minor throat irritations, slightly more energy. Probiotic effects on innate immunity beginning.
Weeks 3-6	Measurable increases in SCFA-producing bacteria. NK cell activity enhanced (shown in de Vrese et al. trial). HLA-DR expression on dendritic cells increasing (Naito et al., 2025). Gut-lung axis becoming more robust.
Weeks 6-12	Full immune priming achieved. If you encounter a cold virus, you should notice shorter duration and milder symptoms compared to previous years. The Shida et al. trial showed URTI incidence dropping from 53% to 22% over 12 weeks.
Full Cold Season	The real payoff: fewer colds overall (24% reduction per Cochrane review), shorter duration when you do catch one, and 42% fewer antibiotic prescriptions. Maintain the protocol for sustained protection.
Year-Round	Continue vitamin D, vitamin C, and prebiotic-rich diet year-round. Resume full protocol 4-6 weeks before next cold season. Your microbiome retains some adaptation between seasons.

Tracking Your Progress

Keep a simple log through cold season to measure the protocol's impact:

- **Colds caught this season vs. previous seasons:** Most adults average 2-4 colds per year. Track each episode (start date, duration, peak severity on a 0-10 scale).
- **Cold duration:** From first symptom to resolution. Compare to your historical average (7-10 days is typical). The evidence suggests 1-2 days shorter with probiotics.
- **Antibiotic use:** Track whether you needed antibiotics for secondary infections. The Cochrane review showed 42% fewer antibiotic prescriptions with probiotics.
- **Protocol compliance:** Did you take your shake, probiotic, and supplements consistently? Missed days compound -- consistency is the strongest predictor of benefit.

For objective measurement, microbiome testing (16S rRNA sequencing) before and after the protocol can confirm increases in Lachnospiraceae, Ruminococcaceae, and Bifidobacterium. Vitamin D levels (25-hydroxyvitamin D) should be tested annually to ensure you are in the optimal 40-60 ng/mL range.

Ready to learn more?

Visit biomeguide.com to explore our probiotic strain database, condition guides, and microbiome testing resources. Every strain mentioned in this protocol has a detailed profile page with links to the original research.

biomeguide.com

Sources

Every recommendation in this protocol is based on published, peer-reviewed clinical research. Below is the complete citation list with DOIs for verification.

1. Zhao Y, Dong BR, Hao Q. (2022) "Probiotics for preventing acute upper respiratory tract infections." Cochrane Database Syst Rev. 8:CD006895. DOI: 10.1002/14651858.CD006895.pub4
2. Li L, Hong K, Sun Q, et al. (2020) "Probiotics for Preventing Upper Respiratory Tract Infections in Adults: A Systematic Review and Meta-Analysis of Randomized Controlled Trials." Evid Based Complement Alternat Med. 2020:8734140. DOI: 10.1155/2020/8734140
3. Shida K, Sato T, Iizuka R, et al. (2017) "Daily intake of fermented milk with Lactobacillus casei strain Shirota reduces the incidence and duration of upper respiratory tract infections in healthy middle-aged office workers." Eur J Nutr. 56(1):45-53. DOI: 10.1007/s00394-015-1056-1
4. Naito T, Morikawa M, Maki A, et al. (2025) "Lactobacillus paracasei strain Shirota suppresses upper respiratory tract infections and activates mononuclear phagocytic cells in healthy Japanese office workers." Biosci Microbiota Food Health. 44(3). DOI: 10.12938/bmfh.2025-004
5. Liu S, Hu P, Du X, et al. (2013) "Lactobacillus rhamnosus GG supplementation for preventing respiratory infections in children: a meta-analysis of randomized, placebo-controlled trials." Indian Pediatr. 50(4):377-381. DOI: 10.1007/s13312-013-0123-z
6. West NP, Horn PL, Pyne DB, et al. (2014) "Probiotic supplementation for respiratory and gastrointestinal illness symptoms in healthy physically active individuals." Clin Nutr. 33(4):581-587. DOI: 10.1016/j.clnu.2013.10.002
7. Turner RB, Lehtoranta L, Hibberd A, et al. (2021) "Effect of Bifidobacterium animalis spp. lactis BI-04 on Rhinovirus-Induced Colds: A Randomized, Placebo-Controlled, Single-Center, Phase II Trial." EClinicalMedicine. 43:101224. DOI: 10.1016/j.eclinm.2021.101224
8. de Vrese M, Winkler P, Rautenberg P, et al. (2006) "Probiotic bacteria reduced duration and severity but not the incidence of common cold episodes in a double blind, randomized, controlled trial." Vaccine. 24(44-46):6670-6674. DOI: 10.1016/j.vaccine.2006.05.048
9. Woodall CA, McGeoch LJ, Hay AD, Hammond A. (2022) "Respiratory tract infections and gut microbiome modifications: A systematic review." PLoS One. 17(1):e0262057. DOI: 10.1371/journal.pone.0262057
10. Chunxi L, Haiyue L, Yanxia L, et al. (2020) "The Gut Microbiota and Respiratory Diseases: New Evidence." J Immunol Res. 2020:2340670. DOI: 10.1155/2020/2340670
11. Chan CKY, Tao J, Chan OS, et al. (2020) "Preventing Respiratory Tract Infections by Synbiotic Interventions: A Systematic Review and Meta-Analysis of Randomized Controlled Trials." Adv Nutr. 11(4):979-988. DOI: 10.1093/advances/nmaa003
12. Williams LM, Stoodley IL, Berthon BS, Wood LG. (2022) "The Effects of Prebiotics, Synbiotics, and Short-Chain Fatty Acids on Respiratory Tract Infections and Immune Function: A Systematic Review and Meta-Analysis." Adv Nutr. 13(1):167-192. DOI: 10.1093/advances/nmab114
13. Lagowska K, Bajerska J. (2021) "Probiotic Supplementation and Respiratory Infection and Immune Function in Athletes: Systematic Review and Meta-Analysis of Randomized Controlled Trials." J Athl Train. 56(11):1213-1223. DOI: 10.4085/592-20
14. Martineau AR, Jolliffe DA, Hooper RL, et al. (2017) "Vitamin D supplementation to prevent acute respiratory tract infections: systematic review and meta-analysis of individual participant data." BMJ. 356:i6583. DOI: 10.1136/bmj.i6583
15. Hemila H, Chalker E. (2013) "Vitamin C for preventing and treating the common cold." Cochrane Database Syst Rev. 2013(1):CD000980. DOI: 10.1002/14651858.CD000980.pub4
16. Nault D, Machingo TA, Shipper AG, et al. (2024) "Zinc for prevention and treatment of the common cold." Cochrane Database Syst Rev. 5:CD014914. DOI: 10.1002/14651858.CD014914.pub2

Disclaimer

This information is for educational purposes only and is not medical advice. Consult your healthcare provider before starting any new supplement regimen, especially if you are pregnant, breastfeeding, taking immunosuppressive medications, or have an autoimmune condition. The common cold is generally self-limiting, but seek medical attention for high fever (>103F/39.4C), difficulty breathing, symptoms lasting more than 10 days, or worsening after initial improvement. Individual results may vary. BiomeGuide.com is not affiliated with any specific supplement manufacturer.

Ready to learn more?

Visit biomeguide.com to explore our probiotic strain database, condition guides, and microbiome testing resources. Every strain mentioned in this protocol has a detailed profile page with links to the original research.

biomeguide.com
