

BIOME GUIDE PROTOCOL

IBS Microbiome Protocol

Evidence-based prebiotic, probiotic, and dietary strategies
to restore gut microbial balance in irritable bowel syndrome

Covers: IBS-D (Diarrhea) | IBS-C (Constipation) | IBS-M (Mixed) | Post-Infectious IBS

Based on 10+ systematic reviews and meta-analyses from PubMed | March 2026

biomeguide.com

What This Protocol Addresses

Irritable bowel syndrome (IBS) is the most common functional gastrointestinal disorder, affecting 10-15% of the global population. It is characterized by chronic abdominal pain, bloating, and altered bowel habits (diarrhea, constipation, or both). While IBS has traditionally been viewed as a disorder of gut-brain interaction, groundbreaking research has revealed a consistent pattern of gut microbial dysbiosis at its core.

This protocol targets the shared microbial imbalance found across all IBS subtypes: depletion of beneficial, butyrate-producing bacteria (particularly *Bifidobacterium*, *Lactobacillus*, *Faecalibacterium prausnitzii*, and *Roseburia*) alongside enrichment of pro-inflammatory species (*E. coli*, *Streptococcus*, *Ruminococcus gnavus*). By restoring this balance through targeted prebiotics, clinically-validated probiotic strains, dietary changes, and evidence-based supplements, you can address the root microbial drivers of your symptoms.

Timeline expectation: Subtle improvements in bloating and stool consistency typically emerge within weeks 2-4. Meaningful improvements in abdominal pain and overall symptom severity are expected by weeks 4-8, based on the duration of the strongest clinical trials. Full microbial remodeling (measurable via microbiome testing) takes 3-6 months of consistent protocol adherence. This is a marathon, not a sprint.

The Gut-IBS Connection

IBS is now understood as a disorder of the microbiota-gut-brain axis. Your gut and brain communicate through neural pathways (the vagus nerve), hormonal signals (over 90% of your body's serotonin is made in the gut), immune mediators (cytokines), and microbial metabolites (short-chain fatty acids like butyrate). When the gut microbiome is disrupted, this communication breaks down. Reduced butyrate production weakens your intestinal barrier, allowing bacterial toxins (LPS) to cross into the bloodstream, triggering low-grade inflammation and visceral hypersensitivity. The result: the pain, bloating, and altered motility that define IBS.

A landmark systematic review and meta-analysis of 23 case-control studies (1,340 participants) confirmed that IBS patients have significantly lower *Bifidobacterium* and *Lactobacillus* levels and significantly higher *E. coli* compared to healthy controls (Wang et al., 2020). A 2024 meta-analysis of 9 studies (1,167 samples) confirmed reduced alpha diversity in IBS patients (Li et al., 2024).

Sources: Wang et al. (2020) J Acad Nutr Diet. DOI: 10.1016/j.jand.2019.05.015; Li et al. (2024) Front Microbiol. DOI: 10.3389/fmicb.2024.1492349

The IBS Microbial Fingerprint

Depleted taxa (targets for restoration):

Bacterium	Role	Evidence
<i>Bifidobacterium</i> spp.	Immune regulation, barrier support, SCFA production	Reduced -1.04 log ₁₀ CFU/g (p<0.01) vs. controls
<i>Lactobacillus</i> spp.	Lactic acid production, serotonin signaling, immune modulation	Reduced -0.57 log ₁₀ CFU/g (p<0.01) vs. controls
<i>Faecalibacterium prausnitzii</i>	Major butyrate producer, anti-inflammatory (inhibits NF-kB)	Reduced; Clostridia class depletion in IBS
<i>Roseburia</i> spp.	Butyrate producer, gut barrier maintenance	Lost relevance as core genus in IBS patients
<i>Blautia</i> spp.	SCFA producer, bile acid metabolism	Significantly reduced in IBS cohorts

Enriched taxa (pro-inflammatory, elevated in IBS):

Bacterium	Concern
<i>Escherichia coli</i>	LPS producer; elevated +0.60 log ₁₀ CFU/g (p<0.01) in IBS
<i>Ruminococcus gnavus</i>	Mucin degrader; consistently increased in IBS-D across multiple studies
<i>Streptococcus</i> spp.	Pro-inflammatory; characteristic of IBS patients
Proteobacteria (Enterobacteriaceae)	LPS-producing phylum; substantial increase in IBS

Sources: Wang et al. (2020); Sanchez-Pellicer et al. (2025) J Clin Med. DOI: 10.3390/jcm14041198; Teige et al. (2024) Clin Exp Gastroenterol. DOI: 10.2147/CEG.S451675

What to AVOID

Processed foods and emulsifiers: Carboxymethylcellulose (CMC) and polysorbate-80, found in many processed foods, directly damage your intestinal mucus layer and promote dysbiosis. Read ingredient labels carefully.

Excessive sugar and refined carbohydrates: Feed pro-inflammatory Proteobacteria and promote small intestinal bacterial overgrowth (SIBO), which overlaps with IBS in up to 78% of cases.

Artificial sweeteners: Sucralose and saccharin alter gut microbiome composition and may worsen symptoms. Stevia appears to be better tolerated.

Excessive alcohol: Increases intestinal permeability ("leaky gut"), disrupts microbiome balance, and can trigger flares in both IBS-D and IBS-C.

Caffeine (IBS-D): Stimulates colonic motility and may worsen diarrhea. If you tolerate it, green tea is preferable for its polyphenol content.

Insoluble fiber (wheat bran): While soluble fiber helps IBS, insoluble fiber from wheat bran can worsen bloating and pain. Choose psyllium, oat fiber, and pectin instead.

Discuss with your doctor: Proton pump inhibitors (PPIs) alter the gut microbiome significantly. If you take PPIs long-term, discuss alternatives or lowest effective dose with your provider. NSAIDs damage the gut barrier; consider safer alternatives when possible.

Dietary Recommendations

Prebiotic-Rich Foods (eat daily)

Onions, garlic, leeks, asparagus are natural sources of inulin and FOS that feed *Bifidobacterium* and *Faecalibacterium*. Start with cooked forms (which are gentler) and small amounts if you are sensitive.

Oats and barley contain beta-glucan, a prebiotic fiber that supports *Bifidobacterium* and produces butyrate. Oats are also a source of soluble fiber beneficial for both IBS-D and IBS-C.

Green bananas and cooled rice/potatoes are rich in resistant starch, which feeds *Roseburia*, *Ruminococcus bromii*, and *Faecalibacterium prausnitzii*. Cooking and cooling starchy foods increases their resistant starch content.

Kiwifruit (2 per day) has been shown in RCTs to improve constipation, increase stool frequency, and soften stool consistency. Especially beneficial for IBS-C.

Polyphenol-Rich Foods (feed *Akkermansia muciniphila*)

Berries (blueberries, cranberries, raspberries), **pomegranate**, **green and black tea**, **dark chocolate** (70%+ cacao), and **extra virgin olive oil** all contain polyphenols that selectively increase *Akkermansia muciniphila*, a keystone species that strengthens your mucus barrier and reduces intestinal permeability. Aim for 2-3 servings of polyphenol-rich foods daily.

IBS Subtype-Specific Guidance

IBS-D: Reduce caffeine intake; limit high-FODMAP fruits initially; consider temporary dairy elimination (lactose intolerance overlaps significantly with IBS-D); emphasize binding foods like white rice, bananas, and oatmeal during flares.

IBS-C: Increase kiwifruit to 2 per day; add prunes (50g/day shown effective in RCTs); ensure adequate hydration (2+ liters daily); psyllium fiber is especially beneficial for IBS-C. Increase physical activity, which promotes motility.

Supplement Protocol

Prebiotic Supplements

These prebiotic fibers feed your depleted beneficial bacteria. **IBS-specific note:** start with HALF doses for the first week and increase gradually. IBS patients are more sensitive to fermentable fibers than the general population.

Prebiotic	Feeds	Product	Dose
GOS	Bifidobacterium (strongest), Lactobacillus	Bimuno GOS powder	3.5g/day (optimal IBS dose per RCT)
Arabinoxylans	Bifidobacterium, Roseburia	Stabilized rice bran (NOW Foods)	1 heaping Tbsp/day
Apple Pectin	Faecalibacterium, Eubacterium, Bifidobacterium	Apple pectin powder	1 heaping Tbsp/day
Resistant Starch	Roseburia, Ruminococcus bromii, Faecalibacterium	Bob's Red Mill Potato Starch or Hi-Maize 260	1 heaping Tbsp/day
Inulin	Bifidobacterium, Faecalibacterium	Pure inulin powder	Start 1/2 tsp, titrate to 1 tsp

Probiotic Recommendations

Unlike some protocols that advise against all probiotics, we recommend specific strains where clinical trial evidence supports their use in IBS. The prebiotic protocol above is the foundation for restoring your endogenous beneficial flora; these probiotics provide additional, evidence-based support.

Strain	Evidence	Dose	Duration
<i>B. longum</i> 35624	RCT (n=362): Significant improvement in pain, bloating, bowel dysfunction, gas (p<0.02). Meta-analysis confirmed. (Whorwell et al., 2006; Yuan et al., 2017)	1x10 ⁸ CFU/day	8+ weeks
<i>L. plantarum</i> 299v	RCT (n=214): 78% rated effect excellent/good vs. 8% placebo (p<0.01). Improved pain severity and frequency. (Ducrotte et al., 2012)	10 billion CFU/day	4-8 weeks
<i>B. coagulans</i> MTCC 5856	RCT (n=36, IBS-D): Reduced bloating, pain, diarrhea (p<0.01). Ranked #1 in network meta-analysis of 43 RCTs. (Majeed et al., 2016; Zhang et al., 2022)	2x10 ⁹ CFU/day	8-12 weeks
<i>B. coagulans</i> Unique IS2	RCT (n=40): Improved pain, bloating, urgency, straining, flatulence, bowel satisfaction. (Gupta & Maity, 2021)	6 billion CFU/day	8-12 weeks

Choosing your strains: You do not need to take all four strains. For general IBS, start with *B. longum* 35624 (sold as Align) and *L. plantarum* 299v (sold as Jarrow Ideal Bowel Support). If you have IBS-D specifically, add *B. coagulans* MTCC 5856 (sold as LactoSpore). **Important:** For *B. longum* 35624, the 10^8 CFU dose is specifically what worked in trials; higher and lower doses did NOT show benefit. More is not better with this strain.

Supporting Supplements

Supplement	Why	Dose	Duration
Peppermint Oil (enteric-coated)	Antispasmodic; RR 2.23 for global symptom improvement in meta-analysis of 9 RCTs (Khanna et al., 2014)	180-225mg, 2-3x/day before meals	4-8 weeks; as needed long-term
Vitamin D3	Meta-analysis of 8 RCTs: significantly reduced IBS severity (SMD -0.77, p=0.04). Modulates microbiome, supports barrier. (Chong et al., 2022)	2,000-5,000 IU/day (test levels)	Ongoing; retest at 3 months
L-Glutamine (IBS-D)	RCT (n=106): 79.6% response vs. 5.8% placebo (p<0.0001). Normalized gut permeability. (Zhou et al., 2019)	5g, 3x daily (15g/day)	8 weeks active; taper to 5g/day
Sodium Butyrate	Direct fuel for colonocytes; supports barrier integrity; anti-inflammatory. RCT showed reduced pain during defecation. (Banasiewicz et al., 2013)	300mg, 2x/day (microencapsulated)	First 60 days
Psyllium Husk (soluble fiber)	Meta-analysis: soluble fiber effective for IBS; normalizes stool form in both IBS-D and IBS-C. (Moayyedi et al., 2014)	Start 3g/day, titrate to 10g/day	Ongoing

Safety notes: Peppermint oil must be enteric-coated to prevent heartburn; avoid if you have severe GERD. L-glutamine should be avoided by those with liver disease or seizure disorders. Test vitamin D levels before supplementing at higher doses; target serum 25(OH)D of 40-60 ng/mL. Always consult your healthcare provider before starting any new supplement, especially if you take prescription medications.

The Daily Protocol

The Daily Prebiotic Shake

The Daily Prebiotic Shake

1. Add at least 12 oz of water to a shaker bottle or blender
2. Add 1 heaping tablespoon of stabilized rice bran (arabinoxylans)
3. Add 1 heaping tablespoon of apple pectin powder
4. Add 1 heaping tablespoon of resistant starch (potato starch or Hi-Maize 260)
5. Add 1 heaping teaspoon of inulin powder (start with 1/2 tsp week 1)
6. Optional: 1/2 teaspoon berry polyphenol powder for *Akkermansia* support
7. Shake well and drink!

Timing: Once daily, morning or evening. IBS-D patients may prefer morning; IBS-C patients may benefit from evening.

Week 1: Use HALF of all prebiotic doses. Increase to full dose by week 2-3 as tolerated.

The Two-Phase Protocol

Phase 1: Active Gut Restoration (Weeks 1-8)

During Phase 1, you are actively rebuilding your beneficial gut flora and reducing inflammation. This phase requires the most consistency.

- Daily prebiotic shake (start half-dose week 1, full dose weeks 2-8)
- Probiotic(s): take daily with or without food (strain-dependent)
- Peppermint oil: 180mg enteric-coated, 2-3x/day, 30 minutes before meals
- L-Glutamine (IBS-D): 5g three times daily, mixed in water
- Sodium butyrate: 300mg twice daily with meals
- Vitamin D3: 5,000 IU/day with a fat-containing meal
- Psyllium: start 3g/day in water, increase by 2g/week to 7-10g/day
- Follow modified low-FODMAP if needed (weeks 1-4), then begin reintroduction
- Avoid processed foods, artificial sweeteners, and excessive alcohol
- GOS prebiotic: 3.5g/day (particularly if following low-FODMAP to preserve *Bifidobacterium*)

Phase 2: Maintenance (Week 9 Onward)

In Phase 2, you transition to a sustainable maintenance routine. The goal is to maintain the microbial gains you achieved in Phase 1.

- Prebiotic shake: reduce to once daily or every other day
 - Continue probiotics (can rotate strains every 2-3 months)
 - Discontinue L-glutamine or reduce to 5g/day
 - Discontinue sodium butyrate (your own bacteria should now produce more)
 - Continue vitamin D3 (retest at 3 months; adjust to maintenance dose)
 - Continue peppermint oil as needed for symptom management
 - Begin structured FODMAP reintroduction (one food group at a time, 3-day intervals)
 - Introduce fermented foods cautiously: start with small amounts of kefir or sauerkraut
 - Continue psyllium fiber at tolerated dose
 - Continue GOS prebiotic if well tolerated
-

What to Expect

Timeframe	What You May Experience
Week 1	Adjustment period. Mild increase in gas or bloating is normal as your microbiome begins to shift. This is a sign the prebiotics are working. The half-dose start helps minimize this.
Weeks 2-4	Bloating and gas typically improve. Stool consistency begins to normalize. Peppermint oil provides noticeable relief from abdominal cramping. L-glutamine (IBS-D) begins reducing stool frequency.
Weeks 4-8	Meaningful improvement in abdominal pain severity and frequency (this is the timeframe of most successful probiotic RCTs). Reduced urgency and improved bowel satisfaction.
Months 2-3	Continued improvement in overall IBS severity. Better stress resilience (gut-brain axis normalization). Improved energy and reduced brain fog as inflammation decreases.
Months 3-6	Full microbial remodeling. Measurable changes on microbiome testing. Transition to maintenance phase. Many patients report feeling like a "different person" at this stage.

Tracking Your Progress

Keeping a simple symptom journal helps you identify what works and provides valuable data if you work with a healthcare provider. Track daily: abdominal pain (0-10 scale), bloating severity, stool form (Bristol Stool Scale), stool frequency, and any food triggers. The IBS Severity Scoring System (IBS-SSS) questionnaire is a validated tool you can use monthly to measure overall progress.

Consider a microbiome test (such as Biomesight, Ombre, or Thryve) at baseline and after 3-6 months to objectively measure changes in your *Bifidobacterium*, *Faecalibacterium*, and overall diversity. This is optional but provides powerful confirmation that the protocol is working at the microbial level.

Learn more at BiomeGuide.com

Every probiotic strain mentioned in this protocol has a detailed profile page on our site with links to the original research. Visit biomeguide.com to explore our probiotic strain database, condition-specific guides, and microbiome testing resources.

Sources

Every recommendation in this protocol is based on published, peer-reviewed clinical research. Below is the complete citation list with DOIs for verification.

1. Soufan F, et al. (2025) "The Gut-Brain Axis in Irritable Bowel Syndrome." Health Science Reports. DOI: 10.1002/hsr2.70660
2. Wang L, et al. (2020) "Gut Microbial Dysbiosis in the Irritable Bowel Syndrome: A Systematic Review and Meta-Analysis." J Acad Nutr Diet, 120(4):565-586. DOI: 10.1016/j.jand.2019.05.015
3. Li X, et al. (2024) "Meta-analysis of gut microbiota alterations in patients with irritable bowel syndrome." Front Microbiol. DOI: 10.3389/fmicb.2024.1492349
4. Sanchez-Pellicer P, et al. (2025) "Comparative Analysis of Gut Microbiota in Patients with IBS." J Clin Med, 14(4):1198. DOI: 10.3390/jcm14041198
5. Teige ES, et al. (2024) "Fecal Bacterial Profile in Patients with IBS." Clin Exp Gastroenterol, 17:123-134. DOI: 10.2147/CEG.S451675
6. Maslennikov R, et al. (2026) "Strain-Specific Systematic Review with Meta-Analysis of Probiotics in IBS." J Clin Med, 15(3):1152. DOI: 10.3390/jcm15031152
7. Goodoory VC, et al. (2023) "Efficacy of Probiotics in IBS: Systematic Review and Meta-analysis." Gastroenterology, 165(5):1206-1218. DOI: 10.1053/j.gastro.2023.07.018
8. Wu Y, et al. (2024) "Efficacy of Probiotics, Prebiotics, Synbiotics, and FMT in IBS." Nutrients, 16(13):2114. DOI: 10.3390/nu16132114
9. Zhang T, et al. (2022) "Efficacy of Probiotics for IBS: A Network Meta-Analysis." Front Cell Infect Microbiol, 12:859967. DOI: 10.3389/fcimb.2022.859967
10. Whorwell PJ, et al. (2006) "Efficacy of Bifidobacterium infantis 35624 in women with IBS." Am J Gastroenterol, 101(7):1581-1590. DOI: 10.1111/j.1572-0241.2006.00734.x
11. Ducrotte P, et al. (2012) "Lactobacillus plantarum 299v improves symptoms of IBS." World J Gastroenterol, 18(30):4012-4018. DOI: 10.3748/wjg.v18.i30.4012
12. Majeed M, et al. (2016) "Bacillus coagulans MTCC 5856 in IBS-D management." Nutrition Journal, 15:21. DOI: 10.1186/s12937-016-0140-6
13. Gupta AK, Maity C. (2021) "Bacillus coagulans LBSC in IBS." Medicine, 100(3):e23641. DOI: 10.1097/MD.00000000000023641
14. Silk DBA, et al. (2009) "GOS prebiotic effects on microbiota and symptoms in IBS." Aliment Pharmacol Ther, 29(5):508-518. DOI: 10.1111/j.1365-2036.2008.03911.x
15. Khanna R, et al. (2014) "Peppermint oil for IBS: systematic review and meta-analysis." J Clin Gastroenterol, 48(6):505-512. DOI: 10.1097/MCG.0b013e3182a88357
16. Chong RIH, et al. (2022) "Vitamin D supplementation for IBS." J Gastroenterol Hepatol, 37(6):993-1003. DOI: 10.1111/jgh.15852
17. Zhou Q, et al. (2019) "Dietary glutamine supplements for postinfectious IBS." Gut, 68(6):996-1002. DOI: 10.1136/gutjnl-2017-315136
18. Banasiewicz T, et al. (2013) "Sodium butyrate reduces abdominal pain in IBS." Colorectal Dis, 15(2):204-209. DOI: 10.1111/j.1463-1318.2012.03152.x
19. Zeraattalab-Motlagh S, et al. (2025) "Nutritional Interventions in IBS: An Umbrella Review." Nutr Rev. DOI: 10.1093/nutrit/nuae107
20. Moayyedi P, et al. (2014) "Fiber supplementation in IBS: systematic review and meta-analysis." Am J Gastroenterol, 109(9):1367-1374. DOI: 10.1038/ajg.2014.195
21. Yuan F, et al. (2017) "Efficacy of B. infantis 35624 in IBS: a meta-analysis." Curr Med Res Opin, 33(7):1191-1197. DOI: 10.1080/03007995.2017.1292230
22. Wilson B, et al. (2020) "B-GOS in Conjunction With Low FODMAP Diet in IBS." Gastroenterology, 159(4):1292-1303. DOI: 10.1053/j.gastro.2020.05.062

This information is for educational purposes only and is not medical advice. Consult your healthcare provider before starting any new supplement regimen, especially if you take prescription medications or have other medical conditions. These protocols are based on published clinical research; individual results may vary. If your symptoms are severe, worsening, or accompanied by red-flag symptoms (unexplained weight loss, blood in stool, fever, or onset after age 50), seek medical evaluation promptly. BiomeGuide.com is not affiliated with any specific supplement manufacturer.

Ready to learn more?

Visit **biomeguide.com** to explore our probiotic strain database, condition guides, and microbiome testing resources. Every strain mentioned in this protocol has a detailed profile page with links to the original research.

biomeguide.com